



Multi-year observations of variable incomplete combustion in the New York megacity

Luke D. Schiferl¹, Cong Cao², Bronte Dalton^{3,4}, Andrew Hallward-Driemeier^{1,5}, Ricardo Toledo-Crow⁶,
and Róisín Commame^{1,5}

¹Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964, USA

²School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY 11794, USA

³Columbia College, Columbia University, New York, NY 10027, USA

⁴Institute for Health Metrics and Evaluation, University of Washington, Seattle, WA 98195, USA

⁵Department of Earth and Environmental Sciences, Columbia University, New York, NY 10027, USA

⁶Advanced Science Research Center, City University of New York, New York, NY 10031, USA

Correspondence: Luke D. Schiferl (schiferl@ldeo.columbia.edu)

Received: 11 January 2024 – Discussion started: 25 January 2024

Revised: 13 June 2024 – Accepted: 8 August 2024 – Published: 13 September 2024

Abstract. Carbon monoxide (CO) is a regulated air pollutant that impacts tropospheric chemistry and is an important indicator of the incomplete combustion of carbon-based fuels. In this study, we used 4 years (2019–2022) of winter and spring (January–May) atmospheric CO observations to quantify and characterize city-scale CO enhancements (ΔCO) from the New York City metropolitan area (NYCMA). We observed large variability in ΔCO , roughly 60 % of which was explained by atmospheric transport from the surrounding surface areas to the measurement sites, with the remaining 40 % due to changes in emissions on sub-monthly timescales. We evaluated the CO emissions from the Emissions Database for Global Atmospheric Research (EDGAR), which has been used to scale greenhouse gas emissions, and found the emissions are much too low in magnitude. During the COVID-19 shutdown in spring 2020, we observed a flattening of the diurnal pattern of CO emissions, consistent with reductions in daytime transportation. Our results highlight the role of meteorology in driving the variability in air pollutants and show that the transportation sector is unlikely to account for the non-shutdown observed CO emission magnitude and variability, an important distinction for determining the sources of combustion emissions in urban regions like the NYCMA.

1 Introduction

Carbon monoxide (CO) is released into the atmosphere from incomplete combustion of carbon-based fuels. Poisonous to humans at high concentrations, CO is a regulated criteria pollutant in many countries including the United States (US), where the exceedance event threshold is an hourly average of 35 ppm (US EPA, 2024). CO emissions come from both anthropogenic (e.g., transportation, manufacturing, power generation) and natural (e.g., wildfires, biomass burning) combustion sources. CO emissions from on-road vehicles in the US have decreased by over 50 % in the past several decades due to improvements in combustion efficiency (Bishop and

Stedman, 2008; Parrish, 2006). These emission reductions have led to lower urban CO concentrations measured in situ at the surface and by satellites throughout the column (Buchholz et al., 2021; Hassler et al., 2016; He et al., 2013; Hedelius et al., 2021; Pommier et al., 2013; Yin et al., 2015). US cities have rarely experienced highly toxic concentrations of CO in recent years, but CO is still a useful tracer for incomplete combustion (e.g., Turnbull et al., 2015). Recent improvements in efficiency may be slowing, however, perhaps due to diminishing returns in catalytic converters (Jiang et al., 2018; McDonald et al., 2013).

CO also impacts atmospheric chemistry by controlling oxidative capacity (OH), which in turn impacts the lifetime

of methane (CH₄) and ozone (O₃) production. CO is therefore also an indirect greenhouse gas. The decreasing trend in the global CO burden has led to a relative increase in CO chemical production, higher methane oxidation by OH, and a shorter CH₄ lifetime (Gaubert et al., 2017).

Given the observed changes to atmospheric CO and their potential implications for chemistry and climate, it is important to consider the timescales over which urban CO emissions are changing and which factors control their variability. The recent decreasing annual trends in CO emissions, such as $\sim 4.6\% \text{ yr}^{-1}$ for the US during 2002–2011 (Yin et al., 2015) and $\sim 4.5\% \text{ yr}^{-1}$ for the Washington, DC–Baltimore region during 2015–2020 (Lopez-Coto et al., 2022), have been attributed to improving vehicle combustion efficiency. Over multiple years of flights Lopez-Coto et al. (2022) also observed consistently lower CO emissions on Sundays, likely from fewer on-road vehicles on these days. On shorter scales, Ren et al. (2018) found significant variability ($\pm 50\%–70\%$ of the mean) in CO emission rates between flights for winters 2015 and 2016 but did not explain the variability. Lopez-Coto et al. (2020) attributed observed variability between several days of flights to the varied sampling of sources that also have high hourly variability, such as power plants and traffic. Hall et al. (2020) found evidence for a vehicle emission dependence on meteorology, where CO emissions relative to nitrogen oxides (NO_x) increased with increasing temperature across the winter-to-summer gradient in the Washington, DC–Baltimore region, likely due to the temperature sensitivity of pollution control equipment on diesel vehicles.

CO inventories track the expected emission changes over time and are frequently used in two ways, despite their uncertainty and limited temporal resolution: (1) as an a priori emission for geostatistical and Bayesian inversion studies which calculate optimal emissions relative to the CO inventory (thereby providing an indirect evaluation of the inventory) and (2) to scale greenhouse gas emissions relative to observed atmospheric ratios. The US Environmental Protection Agency (EPA) National Emissions Inventory (NEI) is often used in inversion studies, is updated every few years for reactive compounds including CO, and has been evaluated substantially: NEI CO emissions are overestimated by up to 2 times compared to observations over multiple inventory years (Brioude et al., 2013; McDonald et al., 2018; Miller et al., 2008; Salmon et al., 2018), although some have found smaller discrepancies (Anderson et al., 2014; Brioude et al., 2011; Castellanos et al., 2011; Gonzalez et al., 2021; Lopez-Coto et al., 2022). The Emissions Database for Global Atmospheric Research (EDGAR), which provides monthly global CO, CH₄, and CO₂ emissions, is less often evaluated, despite being used extensively for scaling greenhouse gas emissions. Methods of greenhouse gas scaling use the observed atmospheric ratio of CO₂ : CO or CH₄ : CO along with the inventory CO emission rate to calculate an observation-derived CO₂ or CH₄ emission rate (Hsu et al., 2010; Wunch et al., 2009). Konovalov et al. (2016) used CO satellite col-

umn measurements to derive scaled CO₂ emissions based on the CO₂ : CO ratio from EDGAR v4.2 (and others). Recently, CO emissions from EDGAR were also used to scale CH₄ emissions from cities across the US based on observed CH₄ : CO ratios from in situ aircraft measurements (v4.3.2; Plant et al., 2019; Ren et al., 2018) and satellite columns (v5.0; Plant et al., 2022). These works generally assume that CO emissions are less uncertain than CH₄ emissions. In one of the few evaluations of EDGAR CO emissions, Ren et al. (2018) found the inventory rate to be within the uncertainty in their mass balance approach. CO emission inventories with a higher spatial and temporal resolution than EDGAR and NEI are also available, but they are regional and sector specific (e.g., for mobile emissions; Gately et al., 2017; McDonald et al., 2014) and are therefore more difficult to evaluate when mixed with other emission sectors in the atmosphere. Long-term in situ CO observations provide an advantage for evaluating seasonal trends in inventories compared to aircraft snapshots and satellite observations but have so far been underutilized.

Restrictions on movement imposed by the COVID-19 pandemic beginning in March 2020 greatly reduced transportation-related emissions for both commuting and leisure activities. These reductions were noted on the global scale for CO₂ (Forster et al., 2020; Le Quéré et al., 2020; Liu et al., 2020) and on the city scale for NO₂ (Goldberg et al., 2020; Shi et al., 2021; Tzortziou et al., 2022), CO (Lopez-Coto et al., 2022; Monteiro et al., 2022), and CO₂ (Monteiro et al., 2022; Turner et al., 2020). City centers also experienced an overall reduction in human activity (with potential implications for building energy and electricity consumption) as suburbanites worked from home during the day and many city dwellers fled to rural areas, but the impact of these population pattern changes on combustion emissions is less clear. The New York City metropolitan area (NYCMA) was the first region in the US impacted by COVID-19 shutdowns. Mobility data indicated a drop of up to 60% in traffic and 90% in public-transit metrics between February and April 2020 (Cao et al., 2023; Forster et al., 2020; Tzortziou et al., 2022). Tzortziou et al. (2022) examined NO₂ changes in the NYCMA and found an average reduction of 32% in the city center during the peak shutdown period (15 March–15 May 2020). Concentrations of volatile organic compounds (VOCs) from both combustion (e.g., benzene) and non-combustion (e.g., personal-care products, industrial chemicals, oxidation products) sources also declined by similar magnitudes during this time (Cao et al., 2023). Residual shutdowns (e.g., school closures) stayed in place until March 2021, with mobility, total column NO₂, and many VOC observations remaining lower than in the pre-shutdown period.

In this study, we analyze multi-year atmospheric CO observations at two sites in the urban core to quantify and characterize the variability in city-scale CO enhancements (ΔCO) from the NYCMA over sub-monthly and sub-daily timescales. We use the observed ΔCO along with an atmo-

spheric transport model to isolate the impacts of meteorology on the observations and evaluate the EDGAR inventory, which is widely used to scale greenhouse gas emissions. We also identify changes to regional CO emissions induced by the peak and residual COVID-19 shutdowns. Using CO as a tracer, this work begins to constrain the uncertainty in urban combustion sources.

2 Methods

2.1 Rooftop observations

Ambient CO dry-mole fractions (units: ppbv, parts per billion by volume) were measured at the Advanced Science Research Center (ASRC) rooftop observatory, a site located 56 m above ground level (93 m above sea level, a.s.l.) in Hamilton Heights, West Harlem, Manhattan (40.81534° N, 73.95033° W) (Fig. S1 in the Supplement). The site samples a mixture of combustion sources including on- and off-road transportation, building energy, manufacturing, and electricity generation. An additional description of the ASRC site is found in Commane et al. (2023), and other in situ observations from this site were used by Cao et al. (2023).

Due to varying availability, several different instruments were used to measure CO over the four subsequent winters and springs (January–May) of the study period (2019–2022). The instruments used in this study are (i) Picarro G2401-m for 2019 and 2020 (reporting at 0.5–1 Hz), (ii) Picarro G2401 for 2021 (reporting at ~ 0.3 Hz), and (iii) Aerodyne SuperDUAL for 2022 (reporting at 1 Hz). Each instrument was calibrated using gas cylinders that are traceable to standards calibrated by the Central Calibration Laboratory (CCL) at the National Oceanographic and Atmospheric Administration (NOAA) Global Monitoring Laboratory (GML) in Boulder, Colorado. CCL maintains the World Meteorological Organization (WMO) CO scale (WMO X2014A). The Aerodyne SuperDUAL setup at ASRC is described by Commane et al. (2023).

We calculate the hourly mean of these CO measurements at the ASRC site for hours with at least 50 % valid sub-hourly measurements (e.g., at least 1800 1 Hz measurements), which are rounded to the nearest 1 ppbv.

2.2 Network observations

We also use hourly CO dry-mole fractions at two EPA Air Quality System (AQS) network sites: (i) the City College of New York (CCNY, a campus of the City University of New York or CUNY) (40.81976° N, 73.94825° W) located in Manhattan 500 m north of the ASRC site and (ii) Cornwall (41.82134° N, 73.29726° W) located in Connecticut ~ 125 km northeast of the other sites (Fig. S1). Both network sites report CO dry-mole fractions at 1 ppbv precision. The CCNY site uses a Teledyne API 300EU analyzer, which has a detection limit of 20 ppbv and is calibrated

weekly. The Cornwall site uses a Thermo Scientific 48i-TLE analyzer, which has a detection limit of 40 ppbv and is auto-calibrated daily. EPA CO observations are calibrated to the National Institute of Standards and Technology (NIST) scale, which is within 5 ppbv of the NOAA–WMO calibration scale at ambient dry-mole fractions (Lee et al., 2017). Neither time-specific uncertainties nor sub-hourly observational variability was reported at either site. The CCNY site is located 45 m a.s.l. and is classified by the EPA AQS as an urban-scale site measuring within 4–50 km, while the Cornwall site is located 505 m a.s.l. and is a regional-scale site sampling air within 50 km to hundreds of kilometers from the site.

2.3 Isolation of city-scale measurements

We use the proximity of the ASRC and CCNY sites (0.5 km apart) to separate the CO measurements representative of the city scale, which we are interested in for characterizing the CO variability in the NYCMA, from those of the local scale, emissions from nearby sources like buildings or roads. Based on the assumption that city-scale observations should be similar at both sites, we define city-scale measurements at each site as any hourly CO observation less than or equal to 3 times the standard deviation (σ) added to the daily mean of the other site. In contrast, local-scale observations have variability greater than the mean + 3σ of the other site. In this categorization, we only use hours when both ASRC and CCNY provide CO observations.

The hourly observed CO mole fractions are highly variable and range from 100 to 3000 ppbv at the ASRC site, while the CCNY site observes fewer high peaks (maximum of 2000 ppbv) (Fig. 1). Our categorization scheme indicates that many of these observed peaks are from local sources near the measurement sites, rather than representative of the broader city scale. As with the observed peaks, these local-scale measurements are more prevalent at ASRC than at CCNY, especially during 2020–2022. A comparison of the city-scale CO observations shows a good correlation (R^2 of 0.61, slope of 0.82) between the sites, consistent with using the 3σ filter (Fig. S2). The local-scale observations are clearly outliers in this relationship, especially the high CO mole fractions at ASRC. We exclude the local-scale observations from the subsequent city-scale analysis. This simple filtering approach is conservative and retains ~ 80 % of the hourly CO measurements at the city scale.

2.4 Background estimation

To characterize and quantify CO from the NYCMA, we must account for the atmospheric CO entering the domain without impact from the study area (“the background”). Following methods developed in Ammoura et al. (2016) and Lopez-Coto et al. (2022), we estimate the hourly background CO at the ASRC and CCNY sites as the 5th percentile of mole




Figure 1. Time series of hourly CO observations used in the analysis categorized as city scale (black) and local scale (red) at the ASRC (left) and CCNY (right) sites for January–June 2019–2022 (top to bottom).

fractions from the previous and following 5 d (10 d total) using only the city-scale CO observations. The background CO mole fractions are only calculated for hours with at least 50 % valid CO data in the 10 d window (i.e., at least 120 valid hours). We tested a range of thresholds from 30 % to 70 % and found little sensitivity in the results to the minimum data availability percentage.

Given its location on the edge of the NYCMA domain and status as a regional measurement site, we use the hourly CO mole fractions from the Cornwall site as an independent estimation of the background CO for the NYCMA domain. The hourly background CO at Cornwall is defined as the mean of the CO observations from the previous and following 5 d (10 d total). As done for the ASRC and CCNY site backgrounds, the Cornwall background CO mole fractions are only calculated for those hours with at least 50 % valid CO data in the 10 d window.

The background CO mole fractions are variable but tend to peak in late winter (Fig. S3). Differences in the background CO calculated from the various sites for a given time (i.e., the uncertainty) may be up to 50 ppbv but are often much lower (~ 5 –10 ppbv). Some CO emitted from the NYCMA may be

sampled at Cornwall on days with strong southwest winds, but the 125 km distance allows for dilution of any plumes. Using Cornwall as a background may lead to a slight underestimate in the magnitude of the observed CO enhancements (Sect. 2.5).

2.5 Calculation of observed CO enhancements

We define the observed CO enhancement (ΔCO) generated by the NYCMA for each hourly city-scale observation as in Eq. (1):

$$\text{observed } \Delta\text{CO} = \text{observed CO} - \text{background CO}, \quad (1)$$

where the observed ΔCO (units: ppbv) is the observed CO dry-mole fraction with the background CO removed. Observed ΔCO is calculated for each of the ASRC and CCNY sites using the observed CO from those sites and both the 5th percentile background for that site and the Cornwall background. To quantify the background uncertainty, we compare the calculated observed ΔCO for hours with valid background CO from both locations.

From the hourly observed ΔCO , we calculate (i) the 10 d mean observed ΔCO centered on each day of the study

period to assess sub-monthly CO variability while removing variability on synoptic timescales and (ii) the mean observed ΔCO for each 2 h period throughout the day (a diurnal pattern) over the weekdays (Monday–Friday) and weekends (Saturday–Sunday) of various periods of at least 30 d to assess sub-daily CO variability. In each case, the mean observed ΔCO is only calculated for periods with at least 50 % valid hourly observed ΔCO . For (i), we tested a range of 4–16 d means and found little sensitivity in the results to the length of the averaging period.

2.6 Emission inventory

We use anthropogenic CO emissions from v6.1 of the Emissions Database for Global Atmospheric Research (EDGAR) inventory for air pollutants, which are available monthly for 2018 at a $0.1^\circ \times 0.1^\circ$ spatial resolution (Crippa et al., 2018, 2020). While not ideal for urban-scale studies such as in the NYCMA, EDGAR provides the best option for linking CO and greenhouse gas emissions. EDGAR CO emissions are greatest in the center of the NYCMA (NYC in Fig. S1), where the combustion from the building energy (12 % of the January–May total), road transportation (43 %), manufacturing (16 %), and power generation (11 %) sectors combine to produce substantial CO emissions for the region. Transportation emissions follow along the road network outward to the rest of the domain. Monthly totals for the study domain show peak emissions in February dropping to the lowest emissions in May, with April close to the annual mean (Table S1). These month-to-month CO emission changes are mostly due to the seasonal pattern in building energy usage (i.e., heating). Monthly variability in the EDGAR CO emissions is also spatially explicit, with a larger absolute change month to month in the NYC core region of the domain, where the absolute emissions are also largest (Fig. S4). However, the outer areas of the domain with lower emission totals experience a greater relative change, since a larger portion of the CO emissions there are from building energy.

Compared to previous versions, EDGAR v6.1 extends activity data through 2018 and updates emission factors for combustion and evaporative sources due to road transportation technology improvement. In the NYCMA, total CO emissions in EDGAR v6.1 decline by 25 % from 2012 to 2018. EDGAR v6.1 CO emissions are also 26 % lower in our domain compared to EDGAR v5.0 for 2015, the latest year available for that version. We do not apply any interannual emission scaling to the EDGAR inventory for our study period due to the high uncertainty in sub-national variability, especially during the COVID-19 shutdown periods, and due to inconsistencies between reported trends in EDGAR and NEI (Plant et al., 2022). EDGAR provides emission uncertainty on a national basis, which most recently is cited as 44 % for US CO emissions in 2012 from EDGAR v4.3.2 (Crippa et al., 2018). While Crippa et al. (2020) suggest methods to implement diurnal variability in EDGAR using

nationwide sector-specific scale factors, we choose to leave emissions constant throughout the day. Instead, we use the observations to define a city-specific diurnal emission scaling (see Sect. 2.9).

CO emissions from biomass burning (i.e., prescribed burning and wildfire) sources are very limited during the winter and spring in the NYCMA, and we do not consider them here. For example, total CO emissions for January–May (for each year in 2012–2018) from the Global Fire Emissions Database v4 with small fires (GFED4s; van der Werf et al., 2017) are less than 1 % of the total EDGAR CO emissions during these months.

2.7 Transport model

The Stochastic Time-Inverted Lagrangian Transport (STILT) model simulates the impact surface fluxes have on the atmospheric mole fraction at a given time and place (Fasoli et al., 2018; Lin et al., 2003). STILT moves idealized particles backward in time with their 3-dimensional movement determined by large-scale winds and random turbulence. Contribution of the surface flux to the atmospheric mole fraction (the surface influence) occurs when any particle resides within the lower half of the planetary boundary layer. The surface influence accumulated by each particle is interpolated to a regular grid based on their locations in time to derive a “surface influence footprint”.

We drive STILT using NOAA High-Resolution Rapid Refresh (HRRR) meteorology (3 km horizontal, hourly temporal resolution) (Benjamin et al., 2016) and refer to the two components together as HRRR–STILT. We configure HRRR–STILT to calculate the surface influence footprint (0.01° horizontal, hourly temporal resolution) for the NYCMA domain (Fig. S1), initiated for each hour of the study period, by running 500 particles backward for 24 h from the ASRC observation site. Using these surface influence footprints, we also calculate the relative NYC surface influence (unitless) by normalizing the summed 24 h footprints for the NYC subdomain.

STILT was previously used by Miller et al. (2008) to assess CO emissions across North America. More recently, STILT has been widely used in anthropogenic greenhouse gas studies in various regions (e.g., Floerchinger et al., 2021; Turner et al., 2016; Lin et al., 2021; Sargent et al., 2018), and the HRRR–STILT coupling has been used in several urban areas (e.g., Turner et al., 2020; Ware et al., 2019). Footprints from the HRRR–STILT setup described here were previously used by Tzortziou et al. (2022), Tao et al. (2022), Wei et al. (2022), and Cao et al. (2023) to investigate NO_2 , ozone, CO_2 , and VOCs, respectively, in the NYCMA.

2.8 Calculation of simulated CO enhancements

We determine the simulated CO enhancement (ΔCO) generated by the NYCMA for each hour of the study period as in




Figure 2. Time series of 10 d mean observed (black, grey) and simulated (red, pink) ΔCO for the New York City metropolitan area (NYCMA) domain at the ASRC (black, red) and CCNY (grey, pink) sites during January–May 2019–2022. Observed ΔCO is plotted as a point at the mean using the two background CO methods, and the vertical range represents the uncertainty in the background CO when both backgrounds are available. ΔCO values are plotted in time at the center of the 10 d averaging period. The COVID-19 shutdown periods are shaded in blue: peak (darker, 15 March–15 May 2020) and residual (lighter, 16 May 2020–15 March 2021).

Eq. (2):

$$\text{simulated } \Delta\text{CO} = \text{inventory CO emission flux} \\ \times \text{surface influence footprint}, \quad (2)$$

where the simulated ΔCO (units: ppbv) is the EDGAR inventory CO emission flux (units: $\text{nmol m}^{-2} \text{s}^{-1}$) multiplied by the 24 h HRRR–STILT surface influence footprint covering the NYCMA (units: $\text{ppbv} (\text{nmol m}^{-2} \text{s}^{-1})^{-1}$). Given the lifetime of CO in the atmosphere (~ 1 month), very limited chemical loss is expected over this 24 h period, so all surface CO emissions intercepted by the footprint will reach the observation site.

Each hour produces a single simulated ΔCO , which we sample to match the valid hourly observed ΔCO for each observation site and background combination. Mean simulated ΔCO is also calculated from the hourly simulated ΔCO as described above for the mean observed ΔCO . To assess the sensitivity of our methods, addressed below, we retain the unsampled simulated ΔCO and also calculate hourly and mean simulated ΔCO using the annual mean EDGAR CO emissions. We also test the sensitivity of simulated ΔCO to the STILT configuration settings impacting mixing (horizontal turbulence, minimum mixing height, and vertical mixing scheme) and to the choice of the meteorological product (NAMS, North American Mesoscale Forecast System at 12 km horizontal resolution; GFS, Global Forecast System at 0.25° ; and GDAS, Global Data Assimilation System at 1°) for a subset of the study period (January–February 2022) that shows high variability in observed ΔCO . The impact of the mixing scheme and meteorological product selection were previously assessed in greenhouse gas studies by Sargent et al. (2021), Hajny et al. (2022), and Tomlin et al. (2023).

2.9 Observed relative emission

Given the lack of sub-monthly and sub-daily variability in the inventory CO emissions from EDGAR, any variability in the mean simulated ΔCO is attributed to variability in the surface influence footprint (i.e., transport meteorology). The remaining variability in mean observed ΔCO that is not captured by the simulated variability is likely then to be due to changes in the CO emissions not included in the inventory. We normalize the observed ΔCO with the simulated ΔCO to calculate the observed relative emission (ORE; unitless) as in Eq. (3), which removes the variability in meteorology and highlights the sensitivity of simulated ΔCO to the observed change in CO emissions for any period:

$$\text{ORE} = \frac{\text{observed } \Delta\text{CO}}{\text{simulated } \Delta\text{CO}}. \quad (3)$$

The ORE value also represents the relative bias in the CO inventory compared to the CO observations, where an ORE value greater than 1 indicates the CO inventory needs to be increased to match the observations, while an ORE value less than 1 indicates the inventory is biased high. Sensitivity experiments (not shown) indicate that on average emission changes suggested by the ORE reach the ASRC site via transport within 2 h, so CO emissions and enhancements can be considered to occur simultaneously even at the shortest diurnal timescale examined here.

3 Results and discussion

We use our observations to quantify the magnitude and variability in the city-scale observed CO enhancements (ΔCO) from the NYC metropolitan area (NYCMA). Then, we use our simulations to infer CO emission variability, identify bias

in the CO emission inventory, and quantify the observed relative emission (ORE) for the region. Finally, we identify the impacts of the COVID-19 shutdowns on CO emissions.

3.1 City-scale observed ΔCO

The observed ΔCO from the NYCMA varies substantially on sub-monthly timescales throughout the winter and spring and across all years of the study (Fig. 2). Mean 10 d observed ΔCO ranges from ~ 75 to ~ 275 ppbv, excluding the 2020 COVID-19 shutdown period (see Sect. 3.3). The winters of 2019, 2020, and 2022 experience extended large peaks (> 100 ppbv) in observed ΔCO with general declines toward spring, while in 2021 the large peak occurs in late March (after COVID-19-related school closures ended), at the beginning of the transition to spring, and observed ΔCO increases again by May. In 2019 and 2020, there is less variability outside of these extended peaks as compared to 2021 and 2022, which show several additional small episodes of more elevated ΔCO (~ 50 ppbv).

Observed ΔCO at the ASRC and CCNY sites are generally consistent and highly correlated, indicating observed ΔCO presented here is representative of city-scale ΔCO . The observed ΔCO at the two sites overlap within the uncertainty presented by the background CO methods, except for slight deviations (~ 25 ppbv) observed in February 2020 and March–April 2022.

There is a clear diurnal pattern in the observed ΔCO , which is largely consistent between the ASRC and CCNY sites (Figs. 3a, S5). Weekday (Monday–Friday) observed ΔCO peaks in the mid-to-late morning (06:00–12:00 EST), falls off in the early afternoon (12:00–15:00 EST), and peaks again in the evening (16:00–21:00 EST). Weekend (Saturday–Sunday) observed ΔCO tends to have a lower peak (or no peak at all) in the late morning. The magnitude and amplitude of the diurnal pattern varies between year and season.

The uncertainty in the observed ΔCO derived from the different background CO calculation methods spans from near 0 to 50 ppbv for the 10 d means, with most uncertainties at ~ 10 –25 ppbv. This uncertainty varies between period and site. For example, the range of background CO is consistently larger for the CCNY site than the ASRC site in 2019, but the CCNY background range is smaller than that of ASRC in 2021. The uncertainty is notably small at both sites during the maximum observed ΔCO of early winter 2020 and throughout the peak COVID-19 shutdown, while the largest uncertainty occurs during May 2022 at the ASRC site. The uncertainty in observed ΔCO associated with the mean diurnal patterns is the same as noted for the 10 d means.




Figure 3. (a) Diurnal time series of mean observed (black) and simulated (red) ΔCO values for the NYCMA domain at the ASRC site. ΔCO are separated by weekday (Monday–Friday, M–F; circles) and weekend (Saturday–Sunday, S–S; triangles) and then averaged every 2 h for the pre-shutdown 2020 (1 January–14 March 2020, top) and peak COVID-19 shutdown (15 March–15 May 2020, bottom) periods. Observed ΔCO is plotted as the mean using the two background CO methods, with the vertical bars representing their range when both backgrounds are available. (b) Diurnal time series of the observed relative emission (ORE), where the ORE value is the ratio of observed ΔCO to simulated ΔCO using EDGAR from (a) for each period. The ORE value is plotted in the same manner as the observed ΔCO in (a).

3.2 Inferred CO emissions and variability

Clearly there is more variability in the observed ΔCO than can be explained by the expected changes in the EDGAR CO emissions alone, which drop only 22 % from February to May. To interpret the variability in the observed ΔCO and to evaluate the inventory, we compare the observed ΔCO from the NYCMA with the simulated ΔCO , which combines emissions with transport meteorology (Fig. 2). The 10 d mean simulated ΔCO is always low (slope of ~ 0.3) in magnitude compared to the observations and shows moderate variability (Fig. 4a–b), except for during and after the 2020 COVID-19 shutdown period (see Sect. 3.3). The simulated ΔCO spans only from 50 to 100 ppbv, resulting in a consistent underestimate of 50 ppbv that can reach 150 ppbv during times of peak observed ΔCO . This strong bias is present regardless of the observation site to which the simulated ΔCO is sampled (or left unsampled) and despite the use of annual mean instead of monthly varying CO emissions (Fig. S6). The underestimation is also larger than the interannual variability in simulated ΔCO to changing meteorological conditions, which suggests the CO emissions from the inventory are the cause of the negative bias, rather than errors in the transport (Fig. S6). Results from sensitivity tests for January–February 2022, a period of highly variable ob-

served ΔCO , further indicate that the uncertainty in transport does not account for the underestimate in simulated ΔCO (Fig. S7). We find that lowering the minimum mixing height in STILT from 250 to 150 m increases simulated ΔCO by only ~ 20 ppbv and changes due to the alternative horizontal turbulence and vertical mixing schemes are nearly zero. All additional meteorological products we tested in STILT are generally consistent with the variability using HRRR but still do not reproduce the large peak in observed ΔCO in February 2022. Driving STILT with NAMS rather than HRRR increases simulated ΔCO by up to ~ 30 ppbv during times of higher surface influence, and GFS and GDAS either match or are ~ 30 ppbv lower than the simulated ΔCO driven by HRRR. Simulated ΔCO is similarly insensitive to the choice of the minimum mixing height and meteorological product (only NAMS shown) on diurnal timescales (Fig. S8).

Excluding the COVID-19 shutdown periods, nearly 60 % (R^2 of 0.52–0.59) of the variability in observed ΔCO is explained by the variability in simulated ΔCO , which is mostly driven by differential meteorology and transport to the ASRC site (Fig. 4a). The similar correlation (R^2 of 0.58–0.61) between the observed ΔCO and the relative NYC surface influence from HRRR–STILT, which is independent from CO emissions, explicitly confirms the impact of transport on observed ΔCO (Fig. 4c). These results are consistent with the comparison at the nearby CCNY site (Fig. 4b, d). Excluding the 2020 peak and residual COVID-19 shutdowns, the distribution of relative NYC surface influence spans a large range that matches the full range of observed ΔCO at the ASRC and CCNY sites (Fig. 4c–d). This comparison indicates that air measured during times of higher observed ΔCO tends to experience more interaction with the NYC surface (and the subdomain's larger emissions). Therefore, the observed ΔCO for the NYCMA is heavily dependent on influence from the NYC subdomain, regardless of CO emission magnitude or variability. For the 10 d mean comparison, the variability in the simulated ΔCO induced using monthly varying rather than mean annual emissions is only 2.4 % outside of the COVID-19 shutdowns (Fig. S9).

The remainder of the variability (~ 40 %) in observed ΔCO is then attributed to actual changes in emissions on sub-monthly timescales not included in the EDGAR CO emission inventory. We can use the observed relative emission (ORE), which is the observed ΔCO normalized by the simulated ΔCO (see Sect. 2.9), to quantify these changes. The ORE value calculated using the 10 d mean indicates the observed CO emissions are 1–3 times the magnitude in the EDGAR inventory, excluding the 2020 peak and residual COVID-19 shutdowns, and as with the observed and simulated ΔCO , the ORE value is quite variable between months and years (Fig. 5). In 2019, the ORE value is above 2 in winter and gradually declines to ~ 1.75 by spring. The winters of 2020 and 2022 experience short-lived peak ORE values of over 2.5, with a return to a high ORE value in spring 2022, although this period is more uncertain given the range of

background CO. The diurnal pattern of ORE shows a clear daytime peak and an even greater range than using the 10 d means, with maximum CO emissions nearly 4 times EDGAR CO for weekdays (weekends are slightly lower) prior to the peak COVID-19 shutdown in 2020 (Fig. 3b). Substantial variability in CO emissions has been observed over short timescales due to variable sampling and is not unexpected given strong diurnal patterns of combustion sources (Lopez-Coto et al., 2020; Ren et al., 2018). The diurnal pattern of emissions observed here could be applied as city-specific scaling factors to CO emissions from all sectors (e.g., Crippa et al., 2020). The diurnal pattern in the ORE value is sensitive to the choice of the meteorological product, which is consistent with difficulties in capturing the timing of the diurnal mixing layer in coastal locations (Fig. S8).

On average, not including the COVID-19 shutdowns, the observed CO emissions for the NYCMA are 95 % (81 %–109 %, varying the background CO) and 90 % (78 %–102 %) higher than in the EDGAR inventory at the ASRC and CCNY sites, respectively. These observed CO emission estimates consistently lie substantially above the EDGAR uncertainty threshold of 44 % and may be more in line with emissions from NEI, which were previously thought to be too high (e.g., Salmon et al., 2018). There is no clear trend in observed CO emissions during our study period between the subsequent winters and springs not impacted by COVID-19 shutdowns, which is inconsistent with the US national trend of CO emission reductions from transportation (e.g., Yin et al., 2015) and suggests other combustion source sectors are a contributing cause of interannual variability.

3.3 COVID-19 impacts on NYCMA CO

The lowest observed ΔCO (~ 50 ppbv) during our study occurs during and after the peak COVID-19 shutdown in March–May 2020 (Fig. 2) and is partly driven by meteorology since the relative NYC surface influence during the shutdown is on the lower half of the overall distribution (Fig. 4c–d). Together, the low observed ΔCO and low relative NYC surface influence during the peak COVID-19 shutdown correspond to a regime that does not occur in any other years. The 2020 COVID-19 shutdowns are also the only period during our study when the magnitude of observed and simulated ΔCO nearly agree (ORE value of ~ 1 , Fig. 5). The large drop in observed CO emissions (38 % at the ASRC site) during the peak COVID-19 shutdown compared to non-shutdown periods is consistent with changes in NO_x emissions from Tzortziou et al. (2022). Matching the observed 60 % reduction in mobility requires a doubling of the EDGAR CO on-road transportation emissions, in which case the modified inventory still underestimates the magnitude of the observed ΔCO peaks in winter.

A higher proportion of variability within the peak COVID-19 shutdown can be attributed to meteorology (85 %–90 %) than other times, which implies relatively constant emis-




Figure 4. (a) Comparison of observed and simulated ΔCO at the ASRC site using ΔCO as in Fig. 2 for various periods surrounding and during the COVID-19 shutdowns: non-shutdown (2019, 16 March–1 June 2021, 2022; black), pre-shutdown 2020 (1 January–14 March 2020; yellow), peak shutdown (15 March–15 May 2020; red), residual shutdown 2020 (16 May–1 June 2020; purple), and residual shutdown 2021 (1 January–15 March 2021; blue). Observed ΔCO is plotted as the mean using the two background CO methods, with the horizontal bars representing their range when both backgrounds are available. For each period, the linear best-fit line, the slope determined by ordinary least squares, the coefficient of determination (R^2), and the number of points considered (N) are shown. The 1 : 1 line is shown in dark grey. (b) Same as in (a) but at the CCNY site. (c) Comparison of observed ΔCO at the ASRC site as in (a) and relative NYC surface influence at the ASRC site for the periods as in (a). For each period, the coefficient of determination (R^2) is shown. (d) Same as in (c) but at the CCNY site.




Figure 5. Time series of the observed relative emission (ORE) for the NYCMA domain at the ASRC (black) and CCNY (grey) sites during January–May 2019–2022, where the ORE value is the ratio of observed ΔCO to simulated ΔCO using EDGAR from Fig. 2. The ORE value is plotted as a point at the mean using the two background CO methods, and the vertical range represents the uncertainty in the background CO when both backgrounds are available. The ORE value is plotted in time at the center of the 10 d averaging period. The COVID-19 shutdown periods are shaded in blue: peak (darker, 15 March–15 May 2020) and residual (lighter, 16 May 2020–15 March 2021).

sions (Fig. 4a–b). Only during the residual shutdown of May 2020 is the simulated ΔCO greater than the observed ΔCO (ORE value of < 1), after the loosening of some restrictions (Figs. 2, 5). Here the CO emissions remain lower than expected by EDGAR, despite the relative rise in observed and simulated ΔCO due to a turn toward greater sampling of the

NYC surface (Fig. 4c–d). This result may be due to the transition away from heating as temperatures increased in May 2020.

We also observe lower CO emissions compared to other years during the less severe residual COVID-19 shutdown in winter 2021, when observed CO emissions are only 59 %

greater than the EDGAR inventory at the ASRC site, a 16 % drop from non-shutdown periods. This result is consistent with lingering restrictions on mobility and reduced observed NO_2 (−30 % compared to 2018–2019). The larger relative reduction in NO_2 , calculated on a monthly basis, compared to the expected emission changes is due to favorable wind conditions, according to Tzortziou et al. (2022). However, the distribution of relative NYC surface influence in our study during the residual shutdown of winter 2021 is not particularly different than during the peak shutdown (Fig. 4c–d). The contrasting outcomes from these studies highlight the importance of considering meteorology and time averaging when connecting emission changes to atmospheric measurements.

There is a clear reduction in magnitude and distinct flattening of the observed and simulated ΔCO diurnal patterns during the COVID-19 shutdowns compared to non-shutdown periods (Figs. 3a, S5). Much of this magnitude change is due to transport differences, and so the ORE value is particularly useful here to see the relative change in emissions (Fig. 3b). The peak COVID-19 shutdown leads to a decrease of $\sim 35\%$ in maximum weekday daytime CO emissions (ORE value drops from 4 to 2.5), likely due to reduced traffic emissions, and extends elevated emissions compared to the overnight minimum an hour earlier and later, indicating a longer rush hour in the city. The weekend daytime maximum emission reduction during the peak COVID-19 shutdown is even greater, nearly 50 %. Only during the overnight hours during this peak shutdown are the observed CO emissions consistent with the EDGAR CO emission magnitude.

4 Conclusions

Continuous in situ observations are useful to quantify and characterize highly variable ΔCO from urban domains such as the NYCMA, especially during emission source transition periods between winter and spring. This observed ΔCO variability is heavily dependent on meteorology and transport, and these factors must be accounted for in air quality studies that try to connect atmospheric observations with emission changes. In the NYCMA, we found a substantial portion of observed ΔCO changes caused by emission variability after removing this weather dependence.

Multiple years of observations capture different seasonal and diurnal patterns in inferred CO emissions that are generally not accounted for in CO emission inventories. More variability is needed in these inventories, which could reasonably be developed and implemented for diurnal patterns using city-specific observations. However, large variability in day-to-day vehicle emissions seems unlikely outside of the weekend effect and extraordinary events such as the COVID-19 pandemic. Therefore, at least in the NYCMA, the combustion source variability on the seasonal and sub-monthly scale lies elsewhere, from sectors such as building heating/cooling

and electricity generation, although the latter is tightly monitored for inefficiencies at large power plants.

The EDGAR CO emissions evaluated here for the NYCMA are greatly underestimated. Studies that scale greenhouse gas emissions by combining observed $\text{CO}_2:\text{CO}$ or $\text{CH}_4:\text{CO}$ with CO inventories could also be underestimating those emissions by 2–3 times. We encourage the evaluation of the emission inventory magnitude itself, rather than only the ratios, for the specific study period and location, since the emission ratios and their trends are uncertain and can vary between inventory year and version.

The COVID-19 shutdowns removed substantial vehicle traffic from the NYCMA, and only then did the inventory match the CO emissions inferred from observations. When the observed drop in transportation during the peak shutdown is reproduced in the inventory, the required increase in CO emissions is still not enough to match the observed ΔCO peaks, which shows that other non-transportation emissions (e.g., stationary sources) are larger than currently accounted for in this inventory. The sources of these unattributed CO emissions must be identified in the NYCMA in order to mitigate carbon fuel combustion for both air quality improvement and, since CO_2 and CH_4 may be co-emitted, to meet greenhouse gas emission targets.

Code and data availability. Data that support the findings of this study are available as listed below. Data for the ASRC rooftop CO observations, NYCMA observed and simulated ΔCO , and relative NYC surface influence are available at <https://doi.org/10.5061/dryad.612jm649n> (Schiferl et al., 2024). Data for the EPA CO observations are available at https://aqs.epa.gov/aqsweb/airdata/download_files.html (EPA AQS, 2023). Data for the EDGAR CO emissions are available at https://edgar.jrc.ec.europa.eu/index.php/dataset_ap61 (EDGAR Team, 2023). Code for the STILT model is available at <https://uataq.github.io/stilt/#/> (Fasoli, 2019). The HRRR, NAMS, GFS, and GDAS Air Resources Laboratory (ARL) files for running STILT are available at <https://www.ready.noaa.gov/archives.php> (ARL, 2024). Analysis and plotting codes in R (<https://www.R-project.org/>, R Core Team, 2020) developed for this study are available from the corresponding author (Luke D. Schiferl, schiferl@ldeo.columbia.edu) upon request. R packages used for analysis and plotting include `ggplot2` (Wickham, 2016), `ggpattern` (<http://github.com/coolbutuseless/ggpattern>, FC, 2021), `magick` (<https://CRAN.R-project.org/package=magick>, Ooms, 2020), `anytime` (<https://CRAN.R-project.org/package=anytime>, Eddelbuettel, 2020), `lubridate` (<https://www.jstatsoft.org/v40/i03/>, Grolemund and Wickham, 2011), `raster` (<https://CRAN.R-project.org/package=raster>, Hijmans, 2020), `ncdf4` (<https://CRAN.R-project.org/package=ncdf4>, Pierce, 2020), and `cowplot` (<https://CRAN.R-project.org/package=cowplot>, Wilke, 2020).

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/acp-24-10129-2024-supplement>.

Author contributions. LDS and RC designed the study. RC and AHD provided calibrated ASRC CO measurements. CC and RTC supported ASRC CO measurements. LDS performed STILT simulations and the primary analysis of the observation–model comparison. RC and BD assisted with the analysis. LDS wrote the paper. All co-authors contributed to the preparation of the manuscript.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. This work was funded by NOAA research grants (grant no. NA20OAR4310306 to Luke D. Schiferl, Róisín Commane, and Cong Cao; grant no. NA21OAR4310235 to Luke D. Schiferl and Róisín Commane). Luke D. Schiferl and Róisín Commane were also funded by support from the Columbia University Department of Earth and Environmental Sciences. The authors thank John Mak for instrumentation and the Advanced Science Research Center (ASRC) for hosting the measurements at the ASRC rooftop observatory. We also thank the STILT development team.

Financial support. This research has been supported by the National Oceanic and Atmospheric Administration (grant nos. NA20OAR4310306 and NA21OAR4310235) and Columbia University.

Review statement. This paper was edited by Christoph Gerbig and reviewed by two anonymous referees.

References

- Ammoura, L., Xueref-Remy, I., Vogel, F., Gros, V., Baudic, A., Bonsang, B., Delmotte, M., Té, Y., and Chevallier, F.: Exploiting stagnant conditions to derive robust emission ratio estimates for CO₂, CO and volatile organic compounds in Paris, *Atmos. Chem. Phys.*, 16, 15653–15664, <https://doi.org/10.5194/acp-16-15653-2016>, 2016.
- Anderson, D. C., Loughner, C. P., Diskin, G., Weinheimer, A., Canty, T. P., Salawitch, R. J., Worden, H. M., Fried, A., Mikoviny, T., Wisthaler, A., and Dickerson, R. R.: Measured and modeled CO and NO_y in DISCOVER-AQ: An evaluation of emissions and chemistry over the eastern US, *Atmos. Environ.*, 96, 78–87, <https://doi.org/10.1016/j.atmosenv.2014.07.004>, 2014.
- ARL: Gridded Meteorological Data Archives, NOAA [data set], <https://www.ready.noaa.gov/archives.php> (last access: 2 January 2024), 2024.
- Benjamin, S. G., Weygandt, S. S., Brown, J. M., Hu, M., Alexander, C. R., Smirnova, T. G., Olson, J. B., James, E. P., Dorell, D. C., Grell, G. A., Lin, H., Peckham, S. E., Smith, T. L., Moninger, W. R., Kenyon, J. S., and Manikin, G. S.: A North American Hourly Assimilation and Model Forecast Cycle: The Rapid Refresh, *Mon. Weather Rev.*, 144, 1669–1694, <https://doi.org/10.1175/MWR-D-15-0242.1>, 2016.
- Bishop, G. A. and Stedman, D. H.: A Decade of On-road Emissions Measurements, *Environ. Sci. Technol.*, 42, 1651–1656, <https://doi.org/10.1021/es702413b>, 2008.
- Brioude, J., Kim, S.-W., Angevine, W. M., Frost, G. J., Lee, S.-H., McKeen, S. A., Trainer, M., Fehsenfeld, F. C., Holloway, J. S., Ryerson, T. B., Williams, E. J., Petron, G., and Fast, J. D.: Top-down estimate of anthropogenic emission inventories and their interannual variability in Houston using a mesoscale inverse modeling technique, *J. Geophys. Res.-Atmos.*, 116, D20305, <https://doi.org/10.1029/2011JD016215>, 2011.
- Brioude, J., Angevine, W. M., Ahmadov, R., Kim, S.-W., Evan, S., McKeen, S. A., Hsie, E.-Y., Frost, G. J., Neuman, J. A., Pollack, I. B., Peischl, J., Ryerson, T. B., Holloway, J., Brown, S. S., Nowak, J. B., Roberts, J. M., Wofsy, S. C., Santoni, G. W., Oda, T., and Trainer, M.: Top-down estimate of surface flux in the Los Angeles Basin using a mesoscale inverse modeling technique: assessing anthropogenic emissions of CO, NO_x and CO₂ and their impacts, *Atmos. Chem. Phys.*, 13, 3661–3677, <https://doi.org/10.5194/acp-13-3661-2013>, 2013.
- Buchholz, R. R., Worden, H. M., Park, M., Francis, G., Deeter, M. N., Edwards, D. P., Emmons, L. K., Gaubert, B., Gille, J., Martínez-Alonso, S., Tang, W., Kumar, R., Drummond, J. R., Clerbaux, C., George, M., Coheur, P.-F., Hurtmans, D., Bowman, K. W., Luo, M., Payne, V. H., Worden, J. R., Chin, M., Levy, R. C., Warner, J., Wei, Z., and Kulawik, S. S.: Air pollution trends measured from Terra: CO and AOD over industrial, fire-prone, and background regions, *Remote Sens. Environ.*, 256, 112275, <https://doi.org/10.1016/j.rse.2020.112275>, 2021.
- Cao, C., Gentner, D. R., Commane, R., Toledo-Crow, R., Schiferl, L. D., and Mak, J. E.: Policy-Related Gains in Urban Air Quality May Be Offset by Increased Emissions in a Warming Climate, *Environ. Sci. Technol.*, 57, 9683–9692, <https://doi.org/10.1021/acs.est.2c05904>, 2023.
- Castellanos, P., Marufu, L. T., Doddridge, B. G., Taubman, B. F., Schwab, J. J., Hains, J. C., Ehrman, S. H., and Dickerson, R. R.: Ozone, oxides of nitrogen, and carbon monoxide during pollution events over the eastern United States: An evaluation of emissions and vertical mixing, *J. Geophys. Res.-Atmos.*, 116, D16307, <https://doi.org/10.1029/2010JD014540>, 2011.
- Commane, R., Hallward-Driemeier, A., and Murray, L. T.: Intercomparison of commercial analyzers for atmospheric ethane and methane observations, *Atmos. Meas. Tech.*, 16, 1431–1441, <https://doi.org/10.5194/amt-16-1431-2023>, 2023.
- Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., van Aardenne, J. A., Monni, S., Doering, U., Olivier, J. G. J., Pagliari, V., and Janssens-Maenhout, G.: Gridded emissions of air pollutants for the period 1970–2012 within EDGAR v4.3.2, *Earth Syst. Sci. Data*, 10, 1987–2013, <https://doi.org/10.5194/essd-10-1987-2018>, 2018.
- Crippa, M., Solazzo, E., Huang, G., Guizzardi, D., Koffi, E., Muntean, M., Schieberle, C., Friedrich, R., and Janssens-Maenhout, G.: High resolution temporal profiles in the Emis-

- sions Database for Global Atmospheric Research, *Sci. Data*, 7, 121, <https://doi.org/10.1038/s41597-020-0462-2>, 2020.
- Eddelbuettel, D.: anytime: Anything to 'POSIXct' or 'Date' Converter, R package version 0.3.9 [code], <https://CRAN.R-project.org/package=anytime> (last access: 17 November 2020), 2020.
- EDGAR Team: Emissions Database for Global Atmospheric Research, Global Air Pollutant Emissions v6.1 [data set], https://edgar.jrc.ec.europa.eu/index.php/dataset_ap61 (last access: 12 April 2023), 2023.
- EPA AQS: AirData Pre-Generated Data Files, Hourly Data for CO [data set], https://aqs.epa.gov/aqsweb/airdata/download_files.html (last access: 4 February 2023), 2023.
- Fasoli, B., Lin, J. C., Bowling, D. R., Mitchell, L., and Mendoza, D.: Simulating atmospheric tracer concentrations for spatially distributed receptors: updates to the Stochastic Time-Inverted Lagrangian Transport model's R interface (STILT-R version 2), *Geosci. Model Dev.*, 11, 2813–2824, <https://doi.org/10.5194/gmd-11-2813-2018>, 2018.
- Fasoli, B.: Stochastic Time-Inverted Lagrangian Transport Model v2.x [code], <https://uataq.github.io/stilt/#/> (last access: 8 November 2019), 2019.
- FC, M.: ggpattern: Geoms with Patterns, GitHub [code], <http://github.com/coolbutuseless/ggpattern> (last access: 7 February 2021), 2021.
- Floerchinger, C., Shepson, P. B., Hajny, K., Daube, B. C., Stirm, B. H., Sweeney, C., and Wofsy, S. C.: Relative flux measurements of biogenic and natural gas-derived methane for seven U.S. cities, *Elementa*, 9, 000119, <https://doi.org/10.1525/elementa.2021.000119>, 2021.
- Forster, P. M., Forster, H. I., Evans, M. J., Gidden, M. J., Jones, C. D., Keller, C. A., Lamboll, R. D., Quéré, C. L., Rogelj, J., Rosen, D., Schleussner, C.-F., Richardson, T. B., Smith, C. J., and Turnock, S. T.: Current and future global climate impacts resulting from COVID-19, *Nat. Clim. Change*, 10, 913–919, <https://doi.org/10.1038/s41558-020-0883-0>, 2020.
- Gately, C. K., Hutyra, L. R., Peterson, S., and Sue Wing, I.: Urban emissions hotspots: Quantifying vehicle congestion and air pollution using mobile phone GPS data, *Environ. Pollut.*, 229, 496–504, <https://doi.org/10.1016/j.envpol.2017.05.091>, 2017.
- Gaubert, B., Worden, H. M., Arellano, A. F. J., Emmons, L. K., Tilmes, S., Barré, J., Martinez Alonso, S., Vitt, F., Anderson, J. L., Alkemade, F., Houweling, S., and Edwards, D. P.: Chemical Feedback From Decreasing Carbon Monoxide Emissions, *Geophys. Res. Lett.*, 44, 9985–9995, <https://doi.org/10.1002/2017GL074987>, 2017.
- Goldberg, D. L., Anenberg, S. C., Griffin, D., McLinden, C. A., Lu, Z., and Streets, D. G.: Disentangling the Impact of the COVID-19 Lockdowns on Urban NO₂ From Natural Variability, *Geophys. Res. Lett.*, 47, e2020GL089269, <https://doi.org/10.1029/2020GL089269>, 2020.
- Gonzalez, A., Millet, D. B., Yu, X., Wells, K. C., Griffis, T. J., Baier, B. C., Campbell, P. C., Choi, Y., DiGangi, J. P., Gvakharia, A., Halliday, H. S., Kort, E. A., McKain, K., Nowak, J. B., and Plant, G.: Fossil Versus Nonfossil CO Sources in the US: New Airborne Constraints From ACT-America and GEM, *Geophys. Res. Lett.*, 48, e2021GL093361, <https://doi.org/10.1029/2021GL093361>, 2021.
- Grolemund, G. and Wickham, H.: Dates and Times Made Easy with lubridate, *J. Stat. Softw.*, 40, 1–25, <https://www.jstatsoft.org/v40/i03/> (last access: 17 November 2020), 2011.
- Hajny, K. D., Floerchinger, C. R., Lopez-Coto, I., Pitt, J. R., Gately, C. K., Gurney, K. R., Hutyra, L. R., Jayarathne, T., Kaeser, R., Roest, G. S., Sargent, M., Stirm, B. H., Tomlin, J., Turner, A. J., Shepson, P. B., and Wofsy, S.: A spatially explicit inventory scaling approach to estimate urban CO₂ emissions, *Elementa*, 10, 00121, <https://doi.org/10.1525/elementa.2021.00121>, 2022.
- Hall, D. L., Anderson, D. C., Martin, C. R., Ren, X., Salawitch, R. J., He, H., Canty, T. P., Hains, J. C., and Dickerson, R. R.: Using near-road observations of CO, NO_y, and CO₂ to investigate emissions from vehicles: Evidence for an impact of ambient temperature and specific humidity, *Atmos. Environ.*, 232, 117558, <https://doi.org/10.1016/j.atmosenv.2020.117558>, 2020.
- Hassler, B., McDonald, B. C., Frost, G. J., Borbon, A., Carslaw, D. C., Civerolo, K., Granier, C., Monks, P. S., Monks, S., Parrish, D. D., Pollack, I. B., Rosenlof, K. H., Ryerson, T. B., von Schneidemesser, E., and Trainer, M.: Analysis of long-term observations of NO_x and CO in megacities and application to constraining emissions inventories, *Geophys. Res. Lett.*, 43, 9920–9930, <https://doi.org/10.1002/2016GL069894>, 2016.
- He, H., Stehr, J. W., Hains, J. C., Krask, D. J., Doddridge, B. G., Vinnikov, K. Y., Canty, T. P., Hosley, K. M., Salawitch, R. J., Worden, H. M., and Dickerson, R. R.: Trends in emissions and concentrations of air pollutants in the lower troposphere in the Baltimore/Washington airshed from 1997 to 2011, *Atmos. Chem. Phys.*, 13, 7859–7874, <https://doi.org/10.5194/acp-13-7859-2013>, 2013.
- Hedelius, J. K., Toon, G. C., Buchholz, R. R., Iraci, L. T., Podolske, J. R., Roehl, C. M., Wennberg, P. O., Worden, H. M., and Wunch, D.: Regional and Urban Column CO Trends and Anomalies as Observed by MOPITT Over 16 Years, *J. Geophys. Res.-Atmos.*, 126, e2020JD033967, <https://doi.org/10.1029/2020JD033967>, 2021.
- Hijmans, R. J.: raster: Geographic Data Analysis and Modeling, R package version 3.4-5 [code], <https://CRAN.R-project.org/package=raster> (last access: 23 December 2020), 2020.
- Hsu, Y.-K., VanCuren, T., Park, S., Jakober, C., Herner, J., FitzGibbon, M., Blake, D. R., and Parrish, D. D.: Methane emissions inventory verification in southern California, *Atmos. Environ.*, 44, 1–7, <https://doi.org/10.1016/j.atmosenv.2009.10.002>, 2010.
- Jiang, Z., McDonald, B. C., Worden, H., Worden, J. R., Miyazaki, K., Qu, Z., Henze, D. K., Jones, D. B. A., Arellano, A. F., Fischer, E. V., Zhu, L., and Boersma, K. F.: Unexpected slowdown of US pollutant emission reduction in the past decade, *P. Natl. Acad. Sci. USA*, 115, 5099–5104, <https://doi.org/10.1073/pnas.1801191115>, 2018.
- Kononov, I. B., Berezin, E. V., Ciais, P., Broquet, G., Zhuravlev, R. V., and Janssens-Maenhout, G.: Estimation of fossil-fuel CO₂ emissions using satellite measurements of “proxy” species, *Atmos. Chem. Phys.*, 16, 13509–13540, <https://doi.org/10.5194/acp-16-13509-2016>, 2016.
- Lee, J., Moon, D., Lee, J., Lim, J., Hall, B., Novelli, P., Brewer, P. J., Miller, M., Murugun, A., Minarro, M. D., Qiao, H., Shuguo, H., Konopelko, L. A., Kustikov, Y. A., Kolobova, A. V., Pankratov, V. V., Wasserman, I. I., Zav'yalov, S. V., Efremova, O. V., Pavlov, M. V., Mitchell, G., Guenther, F., Walden, J., Aoki, N., Shimosaka, T., Tatiana, M., Lagler, F., and Borowiak,

- A.: International comparison CCQM-K84 – carbon monoxide in synthetic air at ambient level, *Metrologia*, 54, 08016, <https://doi.org/10.1088/0026-1394/54/1A/08016>, 2017.
- Le Quéré, C., Jackson, R. B., Jones, M. W., Smith, A. J. P., Abernethy, S., Andrew, R. M., De-Gol, A. J., Willis, D. R., Shan, Y., Canadell, J. G., Friedlingstein, P., Creutzig, F., and Peters, G. P.: Temporary reduction in daily global CO₂ emissions during the COVID-19 forced confinement, *Nat. Clim. Change*, 10, 647–653, <https://doi.org/10.1038/s41558-020-0797-x>, 2020.
- Lin, J. C., Gerbig, C., Wofsy, S. C., Andrews, A. E., Daube, B. C., Davis, K. J., and Grainger, C. A.: A near-field tool for simulating the upstream influence of atmospheric observations: The Stochastic Time-Inverted Lagrangian Transport (STILT) model, *J. Geophys. Res.-Atmos.*, 108, 4493, <https://doi.org/10.1029/2002JD003161>, 2003.
- Lin, J. C., Bares, R., Fasoli, B., Garcia, M., Crosman, E., and Lyman, S.: Declining methane emissions and steady, high leakage rates observed over multiple years in a western US oil/gas production basin, *Sci. Rep.*, 11, 22291, <https://doi.org/10.1038/s41598-021-01721-5>, 2021.
- Liu, Z., Ciais, P., Deng, Z., Lei, R., Davis, S. J., Feng, S., Zheng, B., Cui, D., Dou, X., Zhu, B., Guo, R., Ke, P., Sun, T., Lu, C., He, P., Wang, Y., Yue, X., Wang, Y., Lei, Y., Zhou, H., Cai, Z., Wu, Y., Guo, R., Han, T., Xue, J., Boucher, O., Boucher, E., Chevalier, F., Tanaka, K., Wei, Y., Zhong, H., Kang, C., Zhang, N., Chen, B., Xi, F., Liu, M., Bréon, F.-M., Lu, Y., Zhang, Q., Guan, D., Gong, P., Kammen, D. M., He, K., and Schellnhuber, H. J.: Near-real-time monitoring of global CO₂ emissions reveals the effects of the COVID-19 pandemic, *Nat. Commun.*, 11, 5172, <https://doi.org/10.1038/s41467-020-18922-7>, 2020.
- Lopez-Coto, I., Ren, X., Salmon, O. E., Karion, A., Shepson, P. B., Dickerson, R. R., Stein, A., Prasad, K., and Whetstone, J. R.: Wintertime CO₂, CH₄, and CO Emissions Estimation for the Washington, DC–Baltimore Metropolitan Area Using an Inverse Modeling Technique, *Environ. Sci. Technol.*, 54, 2606–2614, <https://doi.org/10.1021/acs.est.9b06619>, 2020.
- Lopez-Coto, I., Ren, X., Karion, A., McKain, K., Sweeney, C., Dickerson, R. R., McDonald, B. C., Ahn, D. Y., Salawitch, R. J., He, H., Shepson, P. B., and Whetstone, J. R.: Carbon Monoxide Emissions from the Washington, DC, and Baltimore Metropolitan Area: Recent Trend and COVID-19 Anomaly, *Environ. Sci. Technol.*, 56, 2172–2180, <https://doi.org/10.1021/acs.est.1c06288>, 2022.
- McDonald, B. C., Gentner, D. R., Goldstein, A. H., and Harley, R. A.: Long-Term Trends in Motor Vehicle Emissions in U.S. Urban Areas, *Environ. Sci. Technol.*, 47, 10022–10031, <https://doi.org/10.1021/es401034z>, 2013.
- McDonald, B. C., McBride, Z. C., Martin, E. W., and Harley, R. A.: High-resolution mapping of motor vehicle carbon dioxide emissions, *J. Geophys. Res.-Atmos.*, 119, 5283–5298, <https://doi.org/10.1002/2013JD021219>, 2014.
- McDonald, B. C., McKeen, S. A., Cui, Y. Y., Ahmadov, R., Kim, S.-W., Frost, G. J., Pollack, I. B., Peischl, J., Ryerson, T. B., Holloway, J. S., Graus, M., Warneke, C., Gilman, J. B., de Gouw, J. A., Kaiser, J., Keutsch, F. N., Hanisco, T. F., Wolfe, G. M., and Trainer, M.: Modeling Ozone in the Eastern U.S. using a Fuel-Based Mobile Source Emissions Inventory, *Environ. Sci. Technol.*, 52, 7360–7370, <https://doi.org/10.1021/acs.est.8b00778>, 2018.
- Miller, S. M., Matross, D. M., Andrews, A. E., Millet, D. B., Longo, M., Gottlieb, E. W., Hirsch, A. I., Gerbig, C., Lin, J. C., Daube, B. C., Hudman, R. C., Dias, P. L. S., Chow, V. Y., and Wofsy, S. C.: Sources of carbon monoxide and formaldehyde in North America determined from high-resolution atmospheric data, *Atmos. Chem. Phys.*, 8, 7673–7696, <https://doi.org/10.5194/acp-8-7673-2008>, 2008.
- Monteiro, V., Miles, N. L., Richardson, S. J., Turnbull, J., Karion, A., Kim, J., Mitchell, L., Lin, J. C., Sargent, M., Wofsy, S., Vogel, F., and Davis, K. J.: The impact of the COVID-19 lockdown on greenhouse gases: a multi-city analysis of in situ atmospheric observations, *Environ. Res. Commun.*, 4, 041004, <https://doi.org/10.1088/2515-7620/ac66cb>, 2022.
- Ooms, J.: magick: Advanced Graphics and Image-Processing in R, R package version 2.4.0 [code], <https://CRAN.R-project.org/package=magick> (last access: 23 December 2020), 2020.
- Parrish, D. D.: Critical evaluation of US on-road vehicle emission inventories, *Atmos. Environ.*, 40, 2288–2300, <https://doi.org/10.1016/j.atmosenv.2005.11.033>, 2006.
- Pierce, D.: ncd4: Interface to Unidata netCDF (Version 4 or Earlier) Format Data Files, R package version 1.17 [code], <https://CRAN.R-project.org/package=ncdf4> (last access: 17 November 2020), 2020.
- Plant, G., Kort, E. A., Floerchinger, C., Gvakharia, A., Vimont, I., and Sweeney, C.: Large Fugitive Methane Emissions From Urban Centers Along the U.S. East Coast, *Geophys. Res. Lett.*, 46, 8500–8507, <https://doi.org/10.1029/2019GL082635>, 2019.
- Plant, G., Kort, E. A., Murray, L. T., Maasackers, J. D., and Aben, I.: Evaluating urban methane emissions from space using TROPOMI methane and carbon monoxide observations, *Remote Sens. Environ.*, 268, 112756, <https://doi.org/10.1016/j.rse.2021.112756>, 2022.
- Pommier, M., McLinden, C. A., and Deeter, M.: Relative changes in CO emissions over megacities based on observations from space, *Geophys. Res. Lett.*, 40, 3766–3771, <https://doi.org/10.1002/grl.50704>, 2013.
- R Core Team: R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/> (last access: 17 November 2020), 2020.
- Ren, X., Salmon, O. E., Hansford, J. R., Ahn, D., Hall, D., Benish, S. E., Stratton, P. R., He, H., Sahu, S., Grimes, C., Heimburger, A. M. F., Martin, C. R., Cohen, M. D., Stunder, B., Salawitch, R. J., Ehrman, S. H., Shepson, P. B., and Dickerson, R. R.: Methane Emissions From the Baltimore-Washington Area Based on Airborne Observations: Comparison to Emissions Inventories, *J. Geophys. Res.-Atmos.*, 123, 8869–8882, <https://doi.org/10.1029/2018JD028851>, 2018.
- Salmon, O. E., Shepson, P. B., Ren, X., He, H., Hall, D. L., Dickerson, R. R., Stirr, B. H., Brown, S. S., Fibiger, D. L., McDuffie, E. E., Campos, T. L., Gurney, K. R., and Thornton, J. A.: Top-Down Estimates of NO_x and CO Emissions From Washington, D.C.–Baltimore During the WINTER Campaign, *J. Geophys. Res.-Atmos.*, 123, 7705–7724, <https://doi.org/10.1029/2018JD028539>, 2018.
- Sargent, M., Barrera, Y., Nehr Korn, T., Hutyra, L. R., Gately, C. K., Jones, T., McKain, K., Sweeney, C., Hegarty, J., Hardiman, B., Wang, J. A., and Wofsy, S. C.: Anthropogenic and biogenic CO₂

- fluxes in the Boston urban region, *P. Natl. Acad. Sci. USA*, 115, 7491–7496, <https://doi.org/10.1073/pnas.1803715115>, 2018.
- Sargent, M. R., Floerchinger, C., McKain, K., Budney, J., Gottlieb, E. W., Hutyra, L. R., Rudek, J., and Wofsy, S. C.: Majority of US urban natural gas emissions unaccounted for in inventories, *P. Natl. Acad. Sci. USA*, 118, e2105804118, <https://doi.org/10.1073/pnas.2105804118>, 2021.
- Schiferl, L., Commane, R., Cao, C., Hallward-Driemeier, A., and Toledo-Crow, R.: ASRC Rooftop CO observations, NYCMA observed and simulated ΔCO , and relative NYC surface influence, Dryad [data set], <https://doi.org/10.5061/dryad.612jm649n>, 2024.
- Shi, Z., Song, C., Liu, B., Lu, G., Xu, J., Van Vu, T., Elliott, R. J. R., Li, W., Bloss, W. J., and Harrison, R. M.: Abrupt but smaller than expected changes in surface air quality attributable to COVID-19 lockdowns, *Sci. Adv.*, 7, eabd6696, <https://doi.org/10.1126/sciadv.abd6696>, 2021.
- Tao, M., Fiore, A. M., Jin, X., Schiferl, L. D., Commane, R., Judd, L. M., Janz, S., Sullivan, J. T., Miller, P. J., Karambelas, A., Davis, S., Tzortziou, M., Valin, L., Whitehill, A., Civerolo, K., and Tian, Y.: Investigating Changes in Ozone Formation Chemistry during Summertime Pollution Events over the Northeastern United States, *Environ. Sci. Technol.*, 56, 15312–15327, <https://doi.org/10.1021/acs.est.2c02972>, 2022.
- Tomlin, J. M., Lopez-Coto, I., Hajny, K. D., Pitt, J. R., Kaeser, R., Stirm, B. H., Jayarathne, T., Floerchinger, C. R., Commane, R., and Shepson, P. B.: Spatial attribution of aircraft mass balance experiment CO₂ estimations for policy-relevant boundaries: New York City, *Elementa*, 11, 00046, <https://doi.org/10.1525/elementa.2023.00046>, 2023.
- Turnbull, J. C., Sweeney, C., Karion, A., Newberger, T., Lehman, S. J., Tans, P. P., Davis, K. J., Lauvaux, T., Miles, N. L., Richardson, S. J., Cambaliza, M. O., Shepson, P. B., Gurney, K., Patarasuk, R., and Razlivanov, I.: Toward quantification and source sector identification of fossil fuel CO₂ emissions from an urban area: Results from the INFLUX experiment, *J. Geophys. Res.-Atmos.*, 120, 292–312, <https://doi.org/10.1002/2014JD022555>, 2015.
- Turner, A. J., Shusterman, A. A., McDonald, B. C., Teige, V., Harley, R. A., and Cohen, R. C.: Network design for quantifying urban CO₂ emissions: assessing trade-offs between precision and network density, *Atmos. Chem. Phys.*, 16, 13465–13475, <https://doi.org/10.5194/acp-16-13465-2016>, 2016.
- Turner, A. J., Kim, J., Fitzmaurice, H., Newman, C., Worthington, K., Chan, K., Wooldridge, P. J., Köehler, P., Frankenberg, C., and Cohen, R. C.: Observed Impacts of COVID-19 on Urban CO₂ Emissions, *Geophys. Res. Lett.*, 47, e2020GL090037, <https://doi.org/10.1029/2020GL090037>, 2020.
- Tzortziou, M., Kwong, C. F., Goldberg, D., Schiferl, L., Commane, R., Abuhassan, N., Szykman, J. J., and Valin, L. C.: Declines and peaks in NO₂ pollution during the multiple waves of the COVID-19 pandemic in the New York metropolitan area, *Atmos. Chem. Phys.*, 22, 2399–2417, <https://doi.org/10.5194/acp-22-2399-2022>, 2022.
- US EPA: Timeline of Carbon Monoxide (CO) National Ambient Air Quality Standards (NAAQS), US EPA, <https://www.epa.gov/copollution/timeline-carbon-monoxide-co-national-ambient-air-quality-standards-naaqs>, last access: 14 May 2024.
- van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth Syst. Sci. Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.
- Ware, J., Kort, E. A., Duren, R., Mueller, K. L., Verhulst, K., and Yadav, V.: Detecting Urban Emissions Changes and Events With a Near-Real-Time-Capable Inversion System, *J. Geophys. Res.-Atmos.*, 124, 5117–5130, <https://doi.org/10.1029/2018JD029224>, 2019.
- Wei, D., Reinmann, A., Schiferl, L. D., and Commane, R.: High resolution modeling of vegetation reveals large summertime biogenic CO₂ fluxes in New York City, *Environ. Res. Lett.*, 17, 124031, <https://doi.org/10.1088/1748-9326/aca68f>, 2022.
- Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag New York, ISBN: 978-3-319-24277-4, 2016.
- Wilke, C. O.: *cowplot: Streamlined Plot Theme and Plot Annotations for “ggplot2”*, R package version 1.1.0 [code], <https://CRAN.R-project.org/package=cowplot> (last access: 17 November 2020), 2020.
- Wunch, D., Wennberg, P. O., Toon, G. C., Keppel-Aleks, G., and Yavin, Y. G.: Emissions of greenhouse gases from a North American megacity, *Geophys. Res. Lett.*, 36, L15810, <https://doi.org/10.1029/2009GL039825>, 2009.
- Yin, Y., Chevallier, F., Ciais, P., Broquet, G., Fortems-Cheiney, A., Pison, I., and Saunois, M.: Decadal trends in global CO emissions as seen by MOPITT, *Atmos. Chem. Phys.*, 15, 13433–13451, <https://doi.org/10.5194/acp-15-13433-2015>, 2015.